

Publikationen Dr. Franziska Wiebel

- Erdmann R., Wiebel F.F., Flessau A., Rytka J., Beyer A., Fröhlich K.-U., Kunau W.-H.: PAS1, a yeast gene required for peroxisome biogenesis, encodes a member of a novel family of putative ATPases. **Cell** 64: 499-510, 1991.
- Höhfeld J., Mertens D., Wiebel F.F., Kunau W.-H.: Defining components required for assembly in *Saccharomyces cerevisiae*. In "**Membrane Biogenesis and Protein Targeting**", Eds. Neupert W., Lill R., S. 185-207. Elsevier, New York, 1992.
- Wiebel F.F., Kunau W.-H.: The Pas2 protein essential for peroxisome biogenesis is related to ubiquitin-conjugating enzymes. **Nature** 359: 73-76, 1992.
- Kunau W.-H., Beyer A., Franken T., Götte K., Marzioch M., Saidowsky J., Skaletz-Rorowski A., Wiebel F.F.: Two complementary approaches to study peroxisome biogenesis in *Saccharomyces cerevisiae*: Forward and reversed genetics. **Biochimie** 75: 209-224, 1993.
- Wiebel F.F. and Gustafsson J.-Å.: Heterodimeric interaction between retinoid X receptor alpha and orphan nuclear receptor OR1 reveals dimerization-induced activation as a novel mechanism of nuclear receptor activation. **Mol Cell Biol** 17: 3977-3986, 1997.
- Feltkamp, D., Wiebel, F.F., Alberti, S. and Gustafsson, J.-Å., Identification of a novel DNA binding site for nuclear orphan receptor OR1. **J Biol Chem** 274: 10421-10429, 1999.
- Wiebel F.F., Steffensen K.R., Treuter E., Feltkamp D. and Gustafsson J.-Å.: Ligandindependent coregulator recruitment by the triply activatable OR1/retinoid X receptor-alpha nuclear receptor heterodimer. **Mol Endocrinol** 13: 1105-1118, 1999.
- Cesari Weimar F.: Investigating the function of the Ets transcription factor Elk-1 by sitedirected mutagenesis in the mouse genome. **Dissertation** an der Fakultät für Biologie der Universität Tübingen, Institut für Zellbiologie, Abteilung Molekularbiologie (Prof. Dr. A. Nordheim) unter Anleitung von Dr. F. Wiebel. Selbstverlag, 2001.
- Wiebel F.F., Rennekampff V., Vintersten K., Nordheim A.: Generation of mice carrying conditional knockout alleles for the transcription factor SRF. **Genesis** 32: 124-126, 2002.
- Cesari F., Brecht S., Vintersten K., Vuong L.G., Hofmann M., Klingel K., Schnorr J.J., Arsenian S., Schild H., Herdegen T., Wiebel F.F., Nordheim A.: Mice deficient for the ets transcription factor elk-1 show normal immune responses and mildly impaired neuronal gene activation. **Mol Cell Biol** 24: 294-305, 2004.
- Cesari F., Rennekampff V., Vintersten K., Vuong L.G., Seibler J., Bode J., Wiebel F.F., Nordheim A.: Elk-1 knock-out mice engineered by Flp recombinase-mediated cassette exchange. **Genesis** 38: 87-92, 2004.
- Li S., Czubryt M.P., McAnally J., Bassel-Duby R., Richardson J.A., Wiebel F.F., Nordheim A., Olson E.N.: Requirement for serum response factor for skeletal muscle growth and maturation revealed by tissue-specific gene deletion in mice. **Proc Natl Acad Sci U S A** 102: 1082-1087, 2005.
- Alberti S., Krause S.M., Kretz O., Philippar U., Lemberger T., Casanova E., Wiebel F.F., Schwarz H., Frotscher M., Schütz G., Nordheim A.: Neuronal migration in the murine rostral migratory stream requires serum response factor. **Proc Natl Acad Sci U S A** 102: 6148-6153, 2005.